

Endotracheal Suctioning of Mechanically Ventilated Patients With Artificial Airways 2010

An electronic literature search for articles published between January 1990 and October 2009 was conducted by using MEDLINE, CINAHL, and Cochrane Library databases. The update of this clinical practice guideline is the result of reviewing a total of 114 clinical trials, 62 reviews and 6 meta-analyses on endotracheal suctioning. The following recommendations are made following the Grading of Recommendations Assessment, Development, and Evaluation (GRADE) criteria: (1) It is recommended that endotracheal suctioning should be performed only when secretions are present, and not routinely; (2) It is suggested that pre-oxygenation be considered if the patient has a clinically important reduction in oxygen saturation with suctioning; (3) Performing suctioning without disconnecting the patient from the ventilator is suggested; (4) Use of shallow suction is suggested instead of deep suction, based on evidence from infant and pediatric studies; (5) It is suggested that routine use of normal saline instillation prior to endotracheal suction should *not* be performed; (6) The use of closed suction is suggested for adults with high F_IO₂, or PEEP, or at risk for lung derecruitment, and for neonates; (7) Endotracheal suctioning without disconnection (closed system) is suggested in neonates; (8) Avoidance of disconnection and use of lung recruitment maneuvers are suggested if suctioning-induced lung derecruitment occurs in patients with *acute lung injury*; (9) It is suggested that a suction catheter is used that occludes less than 50% the lumen of the endotracheal tube in children and adults, and less than 70% in infants; (10) It is suggested that the duration of the suctioning event be limited to less than 15 seconds. *Key words:* closed suction; endotracheal suction; saline instillation; intratracheal suction; open suction; saline lavage; suction catheter; tracheal suction; clinical practice guideline. [Respir Care 2010;55(6):758–764. © 2010 Daedalus Enterprises]

ETS 1.0 DESCRIPTION

Endotracheal suctioning (ETS) is one of the most common procedures performed in patients with artificial airways. It is a component of bronchial hygiene therapy and mechanical ventilation that involves the mechanical aspiration of pulmonary secretions from a patient's artificial airway to prevent its obstruction.¹ The procedure includes patient preparation, the suctioning event, and follow-up care.

There are 2 methods of endotracheal suctioning based on the selection of catheter: open and closed. The *open* suctioning technique requires disconnecting the patient from the ventilator, while the *closed* suctioning technique involves attachment of a sterile, closed, in-line suction catheter to the ventilator circuit, which allows passage of a suction catheter through the artificial airway without disconnecting the patient from the ventilator. There are also 2 methods of suctioning based on the catheter suction depth selected during the procedure: deep and shallow. *Deep suctioning* is defined as

the insertion of a suction catheter until resistance is met, followed by withdrawal of the catheter by 1 cm before application of negative pressure, and *shallow suctioning* as the insertion of a suction catheter to a predetermined depth, usually the length of the artificial airway plus the adapter.²

ETS 2.0 PATIENT PREPARATION

It is recommended to use smaller catheters whenever possible, since suction pressure seems to have less influence on lung volume loss than catheter size.³ For a given diameter of the endotracheal tube (ETT), the level of negative pressure transmitted to the airway is determined by the combination of the catheter size and the suction pressure. The larger the diameter of the catheter size, the less attenuation of the suction pressure through the airways.⁴

2.1 Diameter of the suction catheter should not exceed one half the inner diameter of the artificial airway in adults, providing an internal-to-external diameter ra-

tio of 0.5 in adults,^{5,6} and 0.5–0.66 in infants and small children.⁷

2.2 In preparation for the suctioning event, delivery of 100% oxygen in pediatric⁸ and adult patients⁹ and 10% increase of baseline in neonates^{10–12} for 30–60 seconds prior to the suctioning event is suggested, especially in patients who are hypoxemic before suctioning.^{13,14} This may be accomplished either:

2.2.1 by adjusting the F_{IO_2} setting on the mechanical ventilator, or

2.2.2 by use of a temporary oxygen-enrichment program available on many microprocessor ventilators.¹⁵

2.2.3 Manual ventilation of the patient is not recommended, as it has been shown to be ineffective for providing delivered F_{IO_2} of 1.0.^{16,17} Practitioners should ensure that PEEP is maintained if no other alternative is available to hyper-oxygenate.

2.3 The negative pressure of the unit must be checked by occluding the end of the suction tubing before attaching it to the suction catheter, and prior to each suctioning event. Suction pressure should be set as low as possible and yet effectively clear secretions. Experimental data to support an appropriate maximum suction level are lacking. Negative pressure of 80–100 mm Hg in neonates¹⁸ and less than 150 mm Hg in adults have been recommended.¹⁹

2.4 The *closed suctioning technique* facilitates continuous mechanical ventilation and oxygenation during the suctioning event.^{20,21}

2.4.1 It may prevent lung derecruitment associated with the use of open-suction system in patients at higher risk of desaturation (eg, premature newborns).^{22–29}

2.4.2 It should be considered in patients requiring high F_{IO_2} and PEEP (eg, acute lung injury).^{30–36}

2.4.3 It neither increases nor decreases the risk of ventilator-associated pneumonia.^{37–39}

2.4.4 Daily changes of in-line suction catheters do not decrease the risk of ventilator-associated pneumonia and is not cost-effective.^{40,41}

2.5 A patient should be placed on a pulse oximeter to assess oxygenation during and following the procedure.

ETS 3.0 PROCEDURE

The suctioning event consists of the placement of a suction catheter through the artificial airway into the trachea and the application of negative pressure as the catheter is being withdrawn. Each pass of the suction catheter into the artificial airway is considered a suctioning event.⁴²

3.1 Shallow suctioning is recommended to prevent trauma to the tracheal mucosa.

3.2 Deep suctioning has not shown superior benefit over shallow suction⁴³ and may be associated with more adverse events.^{44–46}

3.3 The duration of each suctioning event should be no more than 15 seconds.^{8,47,48}

3.4 Sterile technique is encouraged during open suctioning technique.²

3.5 Normal saline instillation. Instillation refers to the administration of aliquots of saline directly into the trachea via an artificial airway. It is hypothesized that normal saline instillation may loosen secretions, increase the amount of secretions removed, and aid in the removal of tenacious secretions. However, there is insufficient evidence to support this hypothesis. Normal saline instillation appears to enhance secretion clearance through cough stimulation in adults,⁴⁹ and a recent report suggests that normal saline instillation prior to suctioning is associated with decreased incidence of ventilator-associated pneumonia in ventilated adult patients.⁵⁰ The great majority of the references used to update this guideline indicate that normal saline instillation is unlikely to be beneficial, and may in fact be harmful.^{17,48,51–53} Therefore, it should not be routinely performed prior to performing endotracheal suctioning.

ETS 4.0 FOLLOW-UP CARE

Following the suctioning event:

4.1 Hyper-oxygenation for at least 1 min by following the same technique(s) used to pre-oxygenate the patient may be used, especially in patients who are hypoxemic before and/or during suctioning.¹⁰

4.2 Hyperventilation should not be routinely used.

4.2.1 Lung-recruitment maneuvers may be attempted in patients with clear evidence of derecruitment.^{30,54,55}

4.3 The patient *should be* monitored for adverse reactions.

ETS 5.0 SETTING

Endotracheal suctioning may be performed by properly trained persons in a wide variety of settings that include (but are not limited to):

5.1 Hospital

5.2 Extended care facility

5.3 Home

5.4 Out-patient clinic

5.5 Physician's office

5.6 Transport vehicle

ETS 6.0 INDICATIONS

- 6.1** The need to maintain the patency and integrity of the artificial airway
- 6.2** The need to remove accumulated pulmonary secretions as evidenced by one of:
 - 6.2.1** sawtooth pattern on the flow-volume loop on the monitor screen of the ventilator and/or the presence of coarse crackles over the trachea are strong indicators of retained pulmonary secretions.^{56,57}
 - 6.2.2** increased peak inspiratory pressure during volume-controlled mechanical ventilation or decreased tidal volume during pressure-controlled ventilation⁵⁸
 - 6.2.3** deterioration of oxygen saturation and/or arterial blood gas values⁵⁸
 - 6.2.4** visible secretions in the airway⁵⁸
 - 6.2.5** patient's inability to generate an effective spontaneous cough
 - 6.2.6** acute respiratory distress⁵⁸
 - 6.2.7** suspected aspiration of gastric or upper-airway secretions
- 6.3** The need to obtain a sputum specimen to rule out or identify pneumonia or other pulmonary infection or for sputum cytology

ETS 7.0 CONTRAINDICATIONS

Endotracheal suctioning is a necessary procedure for patients with artificial airways. Most contraindications are relative to the patient's risk of developing adverse reactions or worsening clinical condition as result of the procedure. When indicated, there is no absolute contraindication to endotracheal suctioning, because the decision to withhold suctioning in order to avoid a possible adverse reaction may, in fact, be lethal.

ETS 8.0 HAZARDS/COMPLICATIONS

- 8.1** Decrease in dynamic lung compliance⁵⁹ and functional residual capacity⁶⁰
- 8.2** Atelectasis^{32,37}
- 8.3** Hypoxia/hypoxemia^{61,62}
- 8.4** Tissue trauma to the tracheal and/or bronchial mucosa⁶³
- 8.5** Bronchoconstriction/bronchospasm^{1,60}
- 8.6** Increased microbial colonization of lower airway^{5,64}
- 8.7** Changes in cerebral blood flow^{65,66} and increased intracranial pressure⁶⁷⁻⁶⁹
- 8.8** Hypertension⁷⁰
- 8.9** Hypotension¹⁷
- 8.10** Cardiac dysrhythmias¹⁷

8.11 Routine use of normal saline instillation may be associated with the following adverse events:

- 8.11.1** excessive coughing⁴⁹
- 8.11.2** decreased oxygen saturation^{53,71-75}
- 8.11.3** bronchospasm
- 8.11.4** dislodgement of the bacterial biofilm that colonizes the ETT into the lower airway^{50,76-78}
- 8.11.5** pain, anxiety, dyspnea^{79,80}
- 8.11.6** tachycardia⁵³
- 8.11.7** increased intracranial pressure^{70,81}

ETS 9.0 LIMITATIONS OF METHOD

Endotracheal suctioning is not a benign procedure, and operators should remain sensitive to possible hazards and complications and take all necessary precautions to ensure patient safety. Secretions in peripheral airways are not and should not be directly removed by endotracheal suctioning.

ETS 10.0 ASSESSMENT OF NEED

Qualified personnel should assess the need for endotracheal suctioning as a routine part of the patient/ventilator system assessment as detailed in section 6.0 Indications.

ETS 11.0 ASSESSMENT OF OUTCOME

- 11.1** Improvement in appearance of ventilator graphics and breath sounds^{57,58}
- 11.2** Decreased peak inspiratory pressure with narrowing of peak inspiratory pressure-plateau pressure; decreased airway resistance or increased dynamic compliance; increased tidal volume delivery during pressure-limited ventilation
- 11.3** Improvement in arterial blood gas values or saturation, as reflected by pulse oximetry (S_{pO_2})
- 11.4** Removal of pulmonary secretions

ETS 12.0 RESOURCES

- 12.1** Necessary Equipment
 - 12.1.1** Vacuum source
 - 12.1.2** Calibrated, adjustable regulator
 - 12.1.3** Collection bottle and connecting tubing
 - 12.1.4** Disposable gloves
 - 12.1.4.1** Sterile (open suction)
 - 12.1.4.2** Clean (closed suction)
 - 12.1.5** Sterile suction catheter
 - 12.1.5.1** For selective main-bronchus suctioning, a curved-tip catheter may be helpful.⁸² The information related to the effectiveness of head turning for selective suctioning is inconclusive.

- 12.1.6 Sterile water and cup (open suction)
- 12.1.7 Goggles, mask, and other appropriate equipment for standard precautions⁸³
- 12.1.8 Oxygen source with a calibrated metering device
- 12.1.9 Pulse oximeter
- 12.1.10 Manual resuscitation bag equipped with an oxygen-enrichment device for emergency backup use
- 12.1.11 Stethoscope

12.2 Optional Equipment

- 12.2.1 Electrocardiograph
- 12.2.2 Sterile sputum trap for culture specimen

12.3 Personnel. Licensed or credentialed respiratory therapists or individuals with similar credentials (eg, MD, RN) who have the necessary training and demonstrated skills to correctly assess need for suctioning, perform the procedure, and adequately evaluate the patient after the procedure.

ETS 13.0 MONITORING

The following should be monitored prior to, during, and after the procedure:

- 13.1 Breath sounds
- 13.2 Oxygen saturation
 - 13.2.1 Skin color
 - 13.2.2 Pulse oximeter
- 13.3 Respiratory rate and pattern
- 13.4 Hemodynamic parameters
 - 13.4.1 Pulse rate
 - 13.4.2 Blood pressure, if indicated and available
 - 13.4.3 Electrocardiogram, if indicated and available
- 13.5 Sputum characteristics
 - 13.5.1 Color
 - 13.5.2 Volume
 - 13.5.3 Consistency
 - 13.5.4 Odor
- 13.6 Cough characteristics
- 13.7 Intracranial pressure, if indicated and available
- 13.8 Ventilator parameters
 - 13.8.1 Peak inspiratory pressure and plateau pressure
 - 13.8.2 Tidal volume
 - 13.8.3 Pressure, flow, and volume graphics, if available
 - 13.8.4 F_{IO_2}

ETS 14.0 FREQUENCY

Although the internal lumen of an ETT decreases substantially after a few days of intubation, due to formation of biofilm,⁸⁴ suctioning should be performed *only* when clin-

ically indicated in order to maintain the patency of the artificial airway used.⁸⁵⁻⁸⁷ Special consideration should be given to the potential complications associated with the procedure.

ETS 15.0 INFECTION CONTROL

15.1 Centers for Disease Control guidelines for standard precautions should be followed.⁸³

15.1.1 If manual ventilation is used, care must be taken not to contaminate the airway.

15.1.2 Sterile technique is encouraged during the entire suctioning event.

15.2 All equipment and supplies should be appropriately disposed of or disinfected.

ETS 16.0 RECOMMENDATIONS

The following recommendations are made following the Grading of Recommendations Assessment, Development, and Evaluation (GRADE)^{88,89} criteria:

16.1 It is recommended that endotracheal suctioning should be performed only when secretions are present, and not routinely. (1C)

16.2 It is suggested that pre-oxygenation be considered if the patient has a clinically important reduction in oxygen saturation with suctioning. (2B)

16.3 Performing suctioning without disconnecting the patient from the ventilator is suggested. (2B)

16.4 Use of shallow suction is suggested instead of deep suction, based on evidence from infant and pediatric studies. (2B)

16.5 It is suggested that routine use of normal saline instillation prior to endotracheal suction should *not* be performed. (2C)

16.6 The use of closed suction is suggested for adults with high F_{IO_2} , or PEEP, or at risk for lung derecruitment (2B), and for neonates (2C).

16.7 Endotracheal suctioning without disconnection (closed system) is suggested in neonates. (2B)

16.8 Avoidance of disconnection and use of lung-recruitment maneuvers are suggested if suctioning-induced lung derecruitment occurs in patients with *acute lung injury*. (2B)

16.9 It is suggested that a suction catheter is used that occludes less than 50% of the lumen of the ETT in children and adults, and less than 70% in infants. (2C)

16.10 It is suggested that the duration of the suctioning event be limited to less than 15 seconds. (2C)

17.0 ETS CPG IDENTIFYING INFORMATION AND AVAILABILITY

17.1 Adaptation

Original Publication: Respir Care 1993;38(5):500-504.

17.2 Guideline Developers

American Association for Respiratory Care Clinical Practice Guidelines Steering Committee

Ruben D Restrepo MD RRT FAARC, Chair

Joel M Brown II RRT

John M Hughes MEd RRT AE-C

17.3 Source(s) of funding

None

17.4 Financial disclosures/conflicts of interest

No conflicts of interest.

17.5 Availability

Interested persons may photocopy these clinical practice guidelines (CPGs) for noncommercial purposes of scientific or educational advancement. Please credit the American Association for Respiratory Care (AARC) and RESPIRATORY CARE. All of the AARC CPGs can be downloaded at no charge at <http://www.rcjournal.com/cpgs>.

REFERENCES

- Guglielminotti J, Desmots J, Dureuil B. Effects of tracheal suctioning on respiratory resistances in mechanically ventilated patients. *Chest* 1998;113(5):1335-1338.
- Koepfel R. Endotracheal tube suctioning in the newborn: a review of the literature. *Newborn Infant Nurs Rev* 2006;6:94-99.
- Copnell B, Dargaville PA, Ryan EM, Kiraly NJ, Chin LO, Mills JF, Tingay DG. The effect of suction method, catheter size and suction pressure on lung volume changes during endotracheal suction in piglets. *Pediatr Res* 2009;66(4):405-410.
- Kiraly NJ, Tingay DG, Mills JF, Morley CJ, Copnell B. Negative tracheal pressure during neonatal endotracheal suction. *Pediatr Res* 2008;64(1):29-33.
- Tiffin NH, Keim MR, Frewen TC. The effects of variations in flow through an insufflating catheter and endotracheal-tube and suction-catheter size on test-lung pressures. *Respir Care* 1990;35(9):889-897.
- Vanner R, Bick E. Tracheal pressures during open suctioning. *Anaesthesia* 2008;63(3):313-315.
- Singh NC, Kisson N, Frewen T, Tiffin N. Physiological responses to endotracheal and oral suctioning in pediatric patients: the influence of endotracheal tube sizes and suction pressures. *Clin Intensive Care* 1991;2(6):345-350.
- Kerem E, Yatsiv I, Goitein KJ. Effect of endotracheal suctioning on arterial blood gases in children. *Intensive Care Med* 1990;16(2):95-99.
- Bourgault AM, Brown CA, Hains SM, Parlow JL. Effects of endotracheal tube suctioning on arterial oxygen tension and heart rate variability. *Biol Res Nurs* 2006;7(4):268-278.
- Pritchard M, Flenady V, Woodgate P. Preoxygenation for tracheal suctioning in intubated, ventilated newborn infants. *Cochrane Database Syst Rev* 2001;(3):CD000427.
- González-Cabello H, Furuya ME, Vargas MH, Tudón H, Garduño J, González-Ayala J. Evaluation of antihypoxic maneuvers before tracheal aspiration in mechanically ventilated newborns. *Pediatr Pulmonol* 2005;39(1):46-50.
- Pritchard MA, Flenady V, Woodgate P. Systematic review of the role of pre-oxygenation for tracheal suctioning in ventilated newborn infants. *J Paediatr Child Health* 2003;39(3):163-165.
- Demir F, Dramali A. Requirement for 100% oxygen before and after closed suction. *J Adv Nurs* 2005;51(3):245-251. Erratum in: *J Adv Nurs* 2005;51(6):660.
- Oh H, Seo W. A meta-analysis of the effects of various interventions in preventing endotracheal suction-induced hypoxemia. *J Clin Nurs* 2003;12(6):912-924.
- Campbell RS, Branson RD. How ventilators provide temporary O₂ enrichment: what happens when you press the 100% suction button? *Respir Care* 1992;37(8):933-937.
- Barnes TA, McGarry WP. Evaluation of ten disposable manual resuscitators. *Respir Care* 1990;35(10):960-968.
- Woodgate PG, Flenady V. Tracheal suctioning without disconnection in intubated ventilated neonates. *Cochrane Database Syst Rev* 2001;(2):CD003065.
- Wilińska M, Zielińska M, Szyreter T, Lesiuk W, Wilkowski J, Ziółkowski J, Swietliński J. [Endotracheal suctioning in neonates and children]. *Med Wieku Rozwoj* 2008;12(4):878-884. *Article in Polish*.
- Plevak D, Ward J. Airway management. In: Burton G, Hodgkin J, editors. *Respiratory care: a guideline to clinical practice*. New York: Lippincott Williams & Wilkins; 1997:555-609.
- Johnson KL, Kearney PA, Johnson SB, Niblett JB, MacMillan NL, McClain RE. Closed versus open endotracheal suctioning: costs and physiologic consequences. *Crit Care Med* 1994;22(4):658-666.
- Lee CK, Ng KS, Tan SG, Ang R. Effect of different endotracheal suctioning systems on cardiorespiratory parameters of ventilated patients. *Ann Acad Med Singapore* 2001;30(3):239-244.
- Woodgate PG, Flenady V. Tracheal suctioning without disconnection in intubated ventilated neonates. *Cochrane Database Syst Rev* 2001;(2):CD003065.
- Lindgren S, Odenstedt H, Olegård C, Söndergaard S, Lundin S, Stenqvist O. Regional lung derecruitment after endotracheal suction during volume- or pressure- controlled ventilation: a study using electric impedance tomography. *Intensive Care Med* 2007;33(1):172-180.
- Kalyn A, Blatz S, Feuerstake S, Paes B, Bautista C. Closed suctioning of intubated neonates maintains better physiologic stability: a randomized trial. *J Perinatol* 2003;23(3):218-222.
- Tan AM, Gomez JM, Mathews J, Williams M, Paratz J, Rajadural VS. Closed versus partially ventilated endotracheal suction in extremely preterm neonates: physiological consequences. *Intensive Crit Care Nurs* 2005;21(4):234-242.
- Cordero L, Sananes M, Ayers LW. Comparison of a closed (Trach Care MAC) with an open endotracheal suction system in small premature infants. *J Perinatol* 2000;20(3):151-156.
- Choong K, Chatkav P, Frndova H, Cox PN. Comparison of loss in lung volume with open versus in-line catheter endotracheal suctioning. *Pediatr Crit Care Med* 2003;4(1):69-73.
- Kiraly NJ, Tingay DG, Mills JF, Morley CJ, Dargaville PA, Copnell B. The effects of closed endotracheal suction on ventilation during conventional and high frequency oscillatory ventilation. *Pediatr Res* 2009;66(4):400-404.
- Hoellner AB, Copnell B, Dargaville PA, Mills JF, Tingay DG. Lung volume and cardiorespiratory changes during open and closed endotracheal suction in ventilated newborn infants. *Arch Dis Child Fetal Neonatal Ed* 2008;93(6):F436-F441.
- Maggiore S, Lellouche F, Pigeot J, Taille S, Deye N, Durrmeyer X, et al. Prevention of endotracheal suctioning-induced alveolar derecruitment in acute lung injury. *Am J Respir Crit Care Med* 2003;167(9):1215-1224.
- Brochard L, Mion G, Isabey D, Bertrand C, Messadi AA, Mancebo J, et al. Constant-flow insufflation prevents arterial oxygen desaturation during endotracheal suctioning. *Am Rev Respir Dis* 1991;144(2):395-400.

32. Lasocki S, Lu Q, Sartorius A, Foulliat D, Remerand F, Rouby JJ. Open and closed- circuit endotracheal suctioning in acute lung injury: efficiency and effects on gas exchange. *Anesthesiology* 2006; 104(1):39-47.
33. Tingay DG, Copnell B, Mills JF, Morley CJ, Dargaville PA. Effects of open endotracheal suction on lung volume in infants receiving HFOV. *Intensive Care Med* 2007;33(4):689-693.
34. Cereda M, Villa F, Colombo E, Greco G, Nacoti M, Presenti A. Closed system endotracheal suctioning maintains lung volume during volume-controlled mechanical ventilation. *Intensive Care Med* 2001;27(4):648-654.
35. Caramaz MP, Schettino G, Suchodolski K, Nishida T, Harris RS, Malhotra A, Kacmarek RM. The impact of endotracheal suctioning on gas exchange and hemodynamics during lung-protective ventilation in acute respiratory distress syndrome. *Respir Care* 2006;51(5): 497-502.
36. Fernández MD, Piacentini E, Blanch L, Fernández R. Changes in lung volume with three systems of endotracheal suctioning with and without pre-oxygenation in patients with mild-to-moderate lung failure. *Intensive Care Med* 2004;30(12):2210-2215.
37. Subirana M, Solà I, Benito S. Closed tracheal suction systems versus open tracheal suction systems for mechanically ventilated adult patients. *Cochrane Database Syst Rev* 2007;(4):CD004581.
38. Topeli A, Harmanci A, Cetinkaya Y, Akdeniz S, Unal S. Comparison of the effect of closed versus open endotracheal suction systems on the development of ventilator-associated pneumonia. *J Hosp Infect* 2004;58(1):14-19.
39. Zeitoun SS, de Barros AL, Diccini S. A prospective, randomized study of ventilator-associated pneumonia in patients using a closed vs. open suction system. *J Clin Nurs* 2003;12(4):484-489.
40. Kollef MH, Prentice D, Shapiro SD, Fraser VJ, Silver P, Trovillon E, et al. Mechanical ventilation with or without daily changes of in-line suction catheters. *Am J Respir Crit Care Med* 1997; 156(2):466-472.
41. Stoller JK, Orens DK, Fatica C, Elliott M, Kester L, Woods J, et al. Weekly versus daily changes of in-line suction catheters: impact on rates of ventilator-associated pneumonia and associated costs. *Respir Care* 2003;48(5):494-499.
42. Gardner DL, Shirland L. Evidence-based guideline for suctioning the intubated neonate and infant. *Neonatal Netw* 2009;28(5):281-302.
43. Spence K, Gillies D, Waterworth L. Deep versus shallow suction of endotracheal tubes in ventilated neonates and young infants. *Cochrane Database Syst Rev* 2003;(3):CD003309.
44. Boothroyd AE, Murthy BV, Darbyshire A, Petros AJ. Endotracheal suctioning causes right upper lobe collapse in intubated children. *Acta Paediatr* 1996;85(12):1422-1425.
45. Ahn Y, Hwang T. The effects of shallow versus deep endotracheal suctioning on the cytological components of respiratory aspirates in high- risk infants. *Respiration* 2003;70(2):172-178.
46. Youngmee A, Yonghoo J. The effects of shallow and deep endotracheal suctioning on oxygen saturation and heart rate in high-risk infants. *Int J Nurs Stud* 2003;40(2):97-104.
47. Pederson CM, Rosendahl-Nielsen M, Hjermand J, Egerod I. Endotracheal suctioning of the adult intubated patient—what is the evidence? *Intensive Crit Care Nurs* 2009;25(1):21-30.
48. Morrow BM, Argent AC. A comprehensive review of pediatric endotracheal suctioning: effects, indications, and clinical practice. *Pediatr Crit Care Med* 2008;9(5):465-477.
49. Gray JE, MacIntyre NR, Kronenberger WG. The effects of bolus normal-saline instillation in conjunction with endotracheal suctioning. *Respir Care* 1990;35(8):785-790.
50. Caruso P, Denari S, Ruiz SA, Demarzo SE, Deheinzelin D. Saline instillation before tracheal suctioning decreases the incidence of ventilator-associated pneumonia. *Crit Care Med* 2009;37(1):32-38.
51. Branson RD. Secretion management in the mechanically ventilated patient. *Respir Care* 2007;52(10):1328-1347.
52. Celik SA, Kanan N. A current conflict: use of isotonic sodium chloride solution on endotracheal suctioning in critically ill patients. *Dimens Crit Care Nurs* 2006;25(1):11-14.
53. Ridling DA, Martin LD, Bratton SL. Endotracheal suctioning with or without instillation of isotonic sodium chloride solution in critically ill children. *Am J Crit Care* 2003;12(3):212-219.
54. Morrow B, Futter M, Argent A. A recruitment maneuver performed after endotracheal suction does not increase dynamic compliance in ventilated pediatric patients: a randomised controlled trial. *Aust J Physiotherapy* 2007;53(3):163-169.
55. Dyhr T, Bonde J, Larsson A. Lung recruitment maneuvers are effective in regaining lung volume and oxygenation after open endotracheal suctioning in acute respiratory distress syndrome. *Crit Care* 2003;7(1):55-62.
56. Guglielminotti J, Alzieu M, Guidet B, Offenstadt G. Bedside detection on retained tracheobronchial secretions in patients receiving mechanical ventilation: is it time for tracheal suction? *Chest* 2000; 118(4):1095-1099.
57. Wood CJ. Can nurses safely assess the need for endotracheal suction in short-term ventilated patients, instead of using routine techniques? *Intensive Crit Care Nurs* 1998;14(4):170-178.
58. Morrow B, Futter M, Argent A. Endotracheal suctioning: from principles to practice. *Intensive Care Med* 2004;30(6):1167-1174.
59. Main E, Castle R, Newham D, Stocks J. Respiratory physiotherapy vs suction: the effects on respiratory function in ventilated infants and children. *Intensive Care Med* 2004;30(6):1144-1151.
60. Heinze H, Sedemund-Adib B, Heringlake M, Gosch UW, Eichler W. Functional residual capacity changes after different endotracheal suctioning methods. *Anesth Analg* 2008;107(3):941-944.
61. Sabirana M, Sola I, Benito S. Closed tracheal suction systems versus open suction systems for mechanically ventilated adult patients. *Cochrane Database Syst Rev* 2007;(4):CD004581.
62. Jongerden IP, Rovers MM, Grypdonck MH, Bonten MJ. Open and closed endotracheal suction systems in mechanically ventilated patients: a meta-analysis. *Crit Care Med* 2007;35(1):260-270.
63. Runtun N. Suctioning artificial airways in children: appropriate technique. *Paediatr Nurs* 1992;18(2):115-118.
64. Freytag CC, Thies FL, König W, Welte T. Prolonged application of closed in-line suction catheters increases microbial colonization of the lower respiratory tract and bacterial growth on catheter surface. *Infection* 2003;31(1):31-37.
65. Skov L, Ryding J, Pryds O, Greisen G. Changes in cerebral oxygenation and cerebral blood volume during endotracheal suctioning in ventilated neonates. *Acta Paediatr* 1992;81(5):389-393.
66. Shah AR, Kurth CD, Gwiazdowski SG, Chance B, Delivoria-Papadopoulos M. Fluctuations in cerebral oxygenation and blood volume during endotracheal suctioning in premature infants. *J Pediatr* 1992;120(5):769-774.
67. Kerr ME, Rudy EB, Weber BB, Stone KS, Turner BS, Orndoff PA, et al. Effect of short-duration hyperventilation during endotracheal suctioning on intracranial pressure in severe head-injured adults. *Nurs Res* 1997;46(4):195-201.
68. Kerr ME, Sereika SM, Orndoff P, Weber B, Rudy EB, Marion D, et al. Effects of neuromuscular blockers and opiates on the cerebrovascular response to endotracheal suctioning in adults with severe head injuries. *Am J Crit Care* 1998;7(3):205-217.
69. Rudy EB, Turner BS, Baun M, Stone KS, Brucia J. Endotracheal suctioning in adults with head injury. *Heart Lung* 1991;20(6):667-674.

70. Evans JC. Reducing the hypoxemia, bradycardia, and apnea associated with suctioning in low birthweight infants. *J Perinatol* 1992; 12(2):137-142.
71. Ackerman M, Gugerty B. The effect of normal saline bolus instillation in artificial airways. *J Soc Otorhinolaryngol Head Neck Nurses* 1990;8:14-17.
72. Reynolds P, Hoffman L, Schlicting R. Effects of normal saline instillation on secretion volume, dynamic compliance and oxygen saturation (abstract). *Am Rev Respir Dis* 1990;141:A574.
73. Kinloch D. Instillation of normal saline during endotracheal suctioning: effects on mixed venous oxygen saturation. *Am J Crit Care* 1999;8(4):231-242.
74. Akgul S, Akyolcu N. Effects of normal saline on endotracheal suctioning. *J Clin Nurs* 2002;11(6):826-830.
75. Ji YR, Kim HS, Park JH. Instillation of normal saline before suctioning in patients with pneumonia. *Yonsei Med J* 2002;43:607-612.
76. Ackerman MH. The effect of saline lavage prior to suctioning. *Am J Crit Care* 1993;2(4):326-330.
77. Hagler DA, Traver GA. Endotracheal saline and suction catheters: sources of lower airway contamination. *Am J Crit Care* 1994;3(6): 444-447.
78. Estes RJ, Meduri GU. The pathogenesis of ventilator-associated pneumonia. I. Mechanisms of bacterial transcolonization and airway inoculation. *Intensive Care Med* 1995;21(4):365-383.
79. Jablonski R. The experience of being mechanically ventilated. *Qual Health Res* 1994;4(2):186-207.
80. O'Neal P, Grap M, Thompson C, Dudley W. Level of dyspnoea experienced in mechanically ventilated adults with and without saline instillation prior to endotracheal suctioning. *Intensive Crit Care Nurs* 2001;17(6):356-363.
81. Chulay M. Why do we keep putting saline down endotracheal tubes? It's time for a change in the way we suction! *Capsules Comments* 1994;2(4):7-11.
82. Kubota Y, Toyoda Y, Kubota H, Asada A, Ueda Y, Hirota Y. Treatment of atelectasis of upper lung lobes. Selective bronchial suctioning with J-shaped catheter tip and guide mark. *Anaesthesia* 1990; 45(10):842-845.
83. Siegel JD, Rhinehart E, Jackson M, Chiarello L; Centers for Disease Control and Prevention, Healthcare Infection Control Practices Advisory Committee. Guidelines for isolation precautions: preventing transmission of infectious agent in health setting 2007. <http://www.cdc.gov/ncidod/dhqp/pdf/guidelines/isolation2007.pdf>. Accessed April 6, 2010.
84. Shah C, Kollef MH. Endotracheal tube intraluminal volume loss among mechanically ventilated patients. *Crit Care Med* 2004;32(1): 120-125.
85. Van de Leur JP, Zwaveling JH, Loef BG, Van der Schans CP. Endotracheal suctioning versus minimally invasive airway suctioning in intubated patients: a prospective randomised controlled trial. *Intensive Care Med* 2003;29(3):426-432. Erratum in: *Intensive Care Med* 2003;29(7):1798.
86. Cordero L, Sananes M, Ayers LW. A comparison of two airway suctioning frequencies in mechanically ventilated, very-low-birth-weight infants. *Respir Care* 2001;46(8):783-788.
87. Wilson G, Hughes G, Rennie J, Morley C. Evaluation of two endotracheal suction regimes in babies ventilated for respiratory distress syndrome. *Early Hum Dev* 1991;25(2):87-90.
88. Guyatt GH, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coello P, Schünemann HJ; GRADE Working Group. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 2008;336(7650):924-926.
89. Jaeschke R, Guyatt GH, Dellinger P, Schünemann H, Levy MM, Kunz R, et al; GRADE Working Group. Use of GRADE grid to reach decisions on clinical practice guidelines when consensus is elusive. *BMJ* 2008;337:a744.